

A test for the search for life on extrasolar planets

Looking for the terrestrial vegetation signature in the Earthshine spectrum

L. Arnold¹, S. Gillet², O. Lardière², P. Riaud^{2,3} and J. Schneider³

¹ Observatoire de Haute-Provence (OHP) CNRS 04870 Saint-Michel-l'Observatoire, France
e-mail: arnold@obs-hp.fr

² Laboratoire d'Interférométrie Stellaire et Exoplanétaire (LISE) CNRS 04870 Saint-Michel-l'Observatoire, France
e-mail: sgohp@obs-hp.fr, lardiere@obs-hp.fr, riaud@obs-hp.fr

³ Observatoire de Paris-Meudon, 92195, Meudon Cedex, France
e-mail: Jean.Schneider@obspm.fr, Pierre.Riaud@obspm.fr

Received 24 December 2001; accepted 6 May 2002

Abstract. We report spectroscopic observations (400–800 nm, $R \approx 100$) of Earthshine in June, July and October 2001 from which normalised Earth albedo spectra have been derived. The resulting spectra clearly show the blue colour of the Earth due to Rayleigh diffusion in its atmosphere. They also show the signatures of oxygen, ozone and water vapour. We tried to extract from these spectra the signature of Earth vegetation. A variable signal (4 to $10 \pm 3\%$) around 700 nm has been measured in the Earth albedo. It is interpreted as being due to the vegetation red edge, expected to be between 2 to 10% of the Earth albedo at 700 nm, depending on models. We discuss the primary goal of the present observations: their application to the detection of vegetation-like biosignatures on extrasolar planets.

Key words. astrobiology – stars: planetary systems

1. Introduction

The search for life on extrasolar planets has become a reasonable goal since the discovery of Earth-mass planets around a pulsar (Wolszczan & Frail 1992) and Jupiter-mass planets around main-sequence stars (Udry & Mayor 2001). Although the detection of Earth-mass planets is not foreseen before space missions (like COROT scheduled for 2004, Schneider *et al.* 1998), it is likely that a significant proportion of main sequence stars have Earth-like companions in their habitable zone. An important question is what type of biosignatures will unveil the possible presence of life on these planets.

Spectral signatures can be of two kinds. A first type consists of biological activity by-products, such as oxygen and its by-product ozone, in association with water vapour, methane and carbon dioxide (Lovelock 1975, Owen 1980, Angel *et al.* 1986). These biogenic molecules present attractive narrow molecular bands. This led in 1993 to the Darwin ESA project (Léger *et al.* 1996), followed by a similar NASA project, Terrestrial Planet Finder (TPF, Angel & Woolf 1997, Beichman *et al.* 1999). But oxygen is not a universal by-product of biological activity as demonstrated

by the existence of anoxygenic photosynthetic bacteria (Blankenship *et al.* 1995).

A second type of biosignature is provided by signs of stellar light transformation into biochemical energy, such as the planet surface colour from vegetation, whatever the bio-chemical details (Labeyrie 1999). This must translate into the planet reflection spectrum by some characteristic spectral features. This signature is necessarily a more robust biomarker than any biogenic gas such as oxygen, since it is a general feature of any photosynthetic activity (here leaving aside chemotrophic biological activity). Unfortunately, it is often not as sharp as single molecular bands: although it is rather sharp for terrestrial vegetation at ≈ 700 nm (Clark 1999, Coliolo *et al.* 2000, see Fig.1), its wavelength structure can vary significantly among bacteria species and plants (Blankenship *et al.* 1995).

Before initiating a search for extrasolar vegetation, it is useful to test if terrestrial vegetation can be detected remotely. This seems possible as long as Earth is observed with a significant spatial resolution (Sagan *et al.* 1993), but is it still the case if Earth is observed as a single dot? A way to observe the Earth as a whole is to observe the Earthshine with the Moon acting like a remote diffuse reflector illuminated by our planet. It has been proposed for some time (Arcichovsky 1912) to look for the vege-




Fig. 1. Reflectance spectra of photosynthetic (green) vegetation, non-photosynthetic (dry) and a soil (from Clark 1999). The so-called vegetation red edge (VRE) is the green vegetation reflectance strong variation from $\approx 5\%$ at 670 nm to $\approx 70\%$ at 800 nm .

tation colour in the Earthshine to use it as a reference for the search of chlorophyll on other planets, but up to now, Earthshine observations apparently did not have sufficient spectral resolution for that purpose (Tikhoff 1914, Danjon 1928, Goode *et al.* 2001). We present in Section 3.1 normalised Earth albedo spectra showing several atmospheric signatures. We show in Section 3.2 how the vegetation signature around 700 nm can be extracted from these spectra.

2. Observations and data reduction

After a first test made in 1999 with the FEROS spectrograph ($R = 48000$) on the La Silla ESO 1.5 m telescope, we have built a dedicated spectrograph mounted on the 80 cm telescope at the Observatoire de Haute-Provence (Table 1).

We have observed the Moon at ascending and descending phases from April to October 2001. Data collected in June, July and October (Table 2) have been of sufficient quality to derive the results described in this article.

Our observation procedure is the following. The long spectrograph slit allows us to record simultaneously the Earthshine and sky background spectra (≈ 80 CCD lines for each). This single exposure is bracketed by two spectra of the Moonlight. Each of the latter is the mean of 10 spectra (totalling $20''$) taken in different regions to smooth the Moon albedo spatial variations. A series of flat fields (tungsten lamp) is recorded just after the previous cycle.

Before getting a final spectrum from binning of CCD lines, a sub-pixel alignment of CCD lines is done to correct the residual angle between the pixel rows and the dispersion direction: Each image is oversampled by a factor of 8 in the dispersion direction. Each line i is then translated to maximize the cross-correlation function from line i with line 1. After the line binning is done, the spectrum is resampled with 1 nm/pixel for convenience.

Table 1. Telescope and spectrograph characteristics.

Parameters	Values
Telescope diameter, f/ratio	80 cm, 16.5
Slit length (unvignetted)	6 arcmin (2 arcmin)
Slit width	1.6 to 7.8 arcsec
Spatial sampling	1.4 arcsec/pixel
Transmission grating	100 lines/mm
Spectral resolution $\lambda/\Delta\lambda$	≈ 50 to 240 at $\lambda = 700\text{ nm}$
Max. spectral range	400 to 900 nm
Spectral sampling	2.6 nm/pixel
CCD Kodak ship	non-ABG KAF-0401E

Table 2. Earthshine observations journal.

Date (yyyy/mm/dd)	Hour (h min UT)	Exposure time (s) ^a	Resolution $\lambda/\Delta\lambda$	Signal to Noise ^b
2001/06/17	02 37 to 03 08	480	120	190
2001/06/18	02 49 to 03 07	360	120	130
2001/06/24	20 20 to 21 03	600	120	210
2001/06/26	20 41 to 21 45	900	120	150
2001/07/23	20 04 to 20 37	240	50	170
2001/07/24	19 54 to 20 56	1080	50	390
2001/07/25	19 57 to 21 24	1440	50	250
2001/10/13	02 16 to 05 00	2640	240	240
2001/10/14	03 35 to 04 04	480	240	100
2001/10/19	17 37 to 17 54	240	240	40
2001/10/21	17 39 to 19 25	1680	240	150

^a Cumulative exposure time from several 120s, 180s or 240s single exposures.

^b Signal to noise ratio at $\lambda = 650\text{ nm}$ for the cumulative spectrum obtained after single images addition and lines binning.

3. Results

3.1. Earth albedo $EA(\lambda)$

Let us define the following spectra: we call the Sun as seen from outside the Earth atmosphere $S(\lambda)$, Earth atmosphere transmittance $AT(\lambda)$, Moonlight $MS(\lambda)$, Earthshine $ES(\lambda)$, Moon albedo $MA(\lambda)$, and Earth albedo $EA(\lambda)$. We have

$$MS(\lambda) = S(\lambda) \times MA(\lambda) \times AT(\lambda) \times g_1, \quad (1)$$

$$ES(\lambda) = S(\lambda) \times EA(\lambda) \times MA(\lambda) \times AT(\lambda) \times g_2. \quad (2)$$

The Earth albedo is simply given by Eq.2/Eq.1, i.e.

$$EA(\lambda) = \frac{ES(\lambda) \times g_1}{MS(\lambda) \times g_2}. \quad (3)$$

Simplifying by $AT(\lambda)$ means that $ES(\lambda)$ and $MS(\lambda)$ should be ideally recorded simultaneously to avoid significant airmass variation and thus Rayleigh scattering bias.




Fig. 2. Examples of measured Earth albedo spectra. Both spectra are normalized to 1 at 600 nm, but the July spectrum is shifted upwards by 0.5 for clarity. The spectral resolution was ≈ 50 in July, and ≈ 240 in October. The July spectrum has been binned to 10 nm/px to mimic the low resolution that might be used for the first extrasolar planet spectrum.

The mean of the two *MS* spectra bracketing $ES(\lambda)$ is thus used to compute $EA(\lambda)$. The g_i terms are geometric factors related to the Sun, Earth and Moon positions. For simplicity, we set g_1 and g_2 equal to 1, equivalent to a spectrum normalization.

Fig.2 shows that the Earth should be seen as a blue object from space. This blue colour has been known for a long time, and has been confirmed by the Apollo astronauts (Kelley 1988). Tikhoff (1912) had already discovered the blue colour of Earthshine and interpreted it as being due to the Rayleigh scattering in the atmosphere. This point will be discussed in more details in Sect.3.2.

The H_2O bands around 690 and 720 nm, and O_2 narrower band at 760 nm are clearly visible with a resolution of $R \approx 50$. The slope variation occurring at ≈ 600 nm is partially the signature of the deepest zone of the broad ozone absorption band (Chappuis band), going from 440 to 760 nm.

3.2. Earth surface reflectance $SR(\lambda)$

To detect the vegetation signature at 700 nm, it is necessary to extract the Earth surface reflectance $SR(\lambda)$, that contains the spectral information on vegetation, from the atmosphere features contained in $AT(\lambda)$. Said differently, it is necessary to remove the atmospheric bands in this spectral region. Surface reflectance $SR(\lambda)$ is usually presented in Earth remote sensing science by a vec-




Fig. 3. An example of a measured 1-airmass atmospheric transmittance, after Rayleigh scattering correction. H_2O and O_2 bands are obviously present, while the broad ozone goes from 450 to 700 nm with maximum absorption at ≈ 600 nm.

tor giving the directional properties of the scattered light (Liang & Strahler 1999, BRDF 2000). But here, we adopt a simpler scalar definition allowing us to write the albedo spectrum $EA(\lambda)$ as the product

$$EA(\lambda) \approx SR(\lambda) \times AT^{\alpha=2}(\lambda) \quad (4)$$

meaning that photons are transmitted once through a one airmass Earth atmosphere, are scattered by the Earth's surface, and then are transmitted back through the atmosphere a second time, giving a power of 2 on $AT(\lambda)$. Clearly α represents an airmass, but its value of 2 is a rough approximation: all photons do not cross twice an airmass of 1, depending on their impact location on Earth, on how they are scattered in the Earth's atmosphere versus their wavelength, and again how the Sun-Earth-Moon triplet is configured (described by a time-dependent vector $\mathbf{g}_3$). Moreover, photons can be reflected by high-altitude clouds having a high albedo, thereby crossing a thinner airmass before going back to space. The latter proportion of photons is also time-dependent, thus implying that $\alpha = \alpha(\mathbf{g}_3, \lambda, t)$ is probably difficult to estimate.

We obtained $AT(\lambda)$ by the ratio of two mean spectra $MS(\lambda)$ taken at two different Moon elevations. This obviously gives only a measure of the local atmosphere transmittance, whereas $AT(\lambda)$ in Eq.4 represents a mean spectrum for the illuminated Earth seen from the Moon.




Fig. 4. An example of Rayleigh correction: The graph shows the 24+26 June spectrum $SR(\lambda)$ (above) after atmospheric absorption correction (Eq.5), but still containing the Rayleigh scattering signature. The spectrum is fitted with a Rayleigh law adjusted over the [500;670nm] window. The fit is then translated (dash) and adjusted to the [740;800nm] region of $SR(\lambda)$ to show the VRE (here $VRE = 7\%$). $SR(\lambda)$ is normalised to 0.3 at 550 nm (Goode *et al.* 2001) to compare to the ocean albedo (McLinden *et al.* 1997). The $SR(\lambda)$ higher slope in the blue is the signature of Rayleigh diffusion in Earth’s atmosphere.

Our measured $AT(\lambda)$ is corrected for Rayleigh scattering and is normalized to 1 airmass (Fig.3). Then Eq.4 gives

$$SR(\lambda) = \frac{EA(\lambda)}{AT^\alpha(\lambda)}. \quad (5)$$

We adjusted α to remove the atmospheric bands in $SR(\lambda)$ between 600 and 800 nm. We found α values between 1 and 3. Most of the time, we treated the O_2 , O_3 and H_2O bands separately with different α to obtain the best correction.

The surface reflectance we obtain in practice from Eq.5 is shown in Fig.4. The albedo spectrum of the ocean is ≈ 0.1 at 500 nm and decreases smoothly to ≈ 0.05 at 750 nm (McLinden *et al.* 1997, Fig.4). But considering these values and the land albedo from Fig.1, associated with a typical cloud cover of 50% for both ocean and land, it can be shown that the higher Earth albedo in the blue in Fig.2 cannot be explained by the higher albedo of the ocean in the blue, but rather by a contribution of Rayleigh diffusion in the atmosphere.

Therefore $SR(\lambda)$ does not represent the pure surface reflectance, but includes uncorrected atmospheric scat-




Fig. 5. An example of data reduction sequence: The graph shows the June albedo spectrum $EA(\lambda)$ (bottom). All atmospheric absorption features are then corrected according to Eq.5 and the spectrum is flattened with a Rayleigh law adjusted in the [500;670nm] window (Fig.4). The result is shown above with 1 and 10 nm/px resolution. The measured red edge around 700nm is $VRE = 7\%$ (Eq.6).

tering (for simplicity, we nevertheless continue to name $SR(\lambda)$ the result of Eq.5). The Fig.4 shows $SR(\lambda)$ fitted with the Rayleigh law $A + B/\lambda^4$ adjusted over the [500;670nm] window. The slope towards the blue does not hide the relatively sharp vegetation signature, which appears around 700 nm. $SR(\lambda)$ is then normalized to the Rayleigh fit (Fig.5).

To quantify the vegetation signature, we define the Vegetation Red Edge (VRE) as

$$VRE = \frac{r_I - r_R}{r_R} \quad (6)$$

where r_R and r_I are the mean reflectances in the [600;670nm] and [740;800nm] windows in the spectrum after it has been flattened with a Rayleigh law as explained above. This VRE definition giving the relative height of the step due to the vegetation is close to the NDVI (Normalized Difference Vegetation Index, Rouse *et al.* 1974, Tucker 1979) used in Earth satellite observation which considers the difference, after atmospheric correction, between the reflected fluxes in broad red and infra-red bands, normalized to the sum of the fluxes in these bands,

$$NDVI = \frac{f_I - f_R}{f_R + f_I}. \quad (7)$$

Flattened $SR(\lambda)$ spectra are shown in Fig.6 and 7. Eq.6 gives VRE values ranging between 4 and 10%.




Fig. 6. Collection of $SR(\lambda)$ spectra normalized to 1 at 600 nm, but shifted upwards for clarity by 0.2, 0.4, 0.6 and 0.8, respectively. Note that only the [600;670] and [740;800 nm] windows are used to estimate the VRE.




Fig. 7. The same collection of $SR(\lambda)$ spectra as in Fig. 6 but binned to 10 nm/px.

4. Discussion

First, we verified that the Moon albedo cancels correctly in practice when we do the ratio $EA(\lambda) = ES(\lambda)/MS(\lambda)$ (Eq. 3). Since $MS(\lambda)$ is the mean of 10 long-slit Moonlight spectra totalling 20' while a single Earthshine $ES(\lambda)$ spectra is only 2', we computed the ratio of 2 spectra (2'/20')

Table 3. Results for the measured Vegetation Red Edge VRE.

Epoch	Viewed zone	VRE
2001/06/17&18 morning	Eur./Asia	$10 \pm 5\%$
2001/06/24&26 evening	Amer./Pacific	$7 \pm 3\%$
2001/07/23&24&25 evening	Amer./Pacific	$7 \pm 3\%$
2001/10/13&14 morning	Eur./Asia	$4 \pm 3\%$
2001/10/19&21 evening	Amer./Pacific	$7 \pm 3\%$

of Moonlight $MS(\lambda)$ taken on different Moon regions. We obtain a constant with $\sigma < 0.5\%$ meaning that the Moon albedo cancels correctly in Eq. 3.

We also verified that the second order spectrum pollution is negligible: it has been measured to be 0% at 760 nm, < 0.3% at 780 nm and < 0.5% at 800 nm.

To test our VRE measurements, we also measured the VRE on spectra of Vega and the sunlit Moon, for which we obviously should have $VRE = 0\%$. After standard bias, dark and flat corrections, the data are calibrated with reference A0V and Sun spectra, respectively, then flattened by normalization to a black body curve. Vega spectra show a $VRE = -1\%$, $\sigma = 2\%$. We also used the standard Moon albedo spectrum of Apollo16 sample 62231 (Pieters 1999) to properly flatten the Moon data: We obtain $VRE = 0\%$, $\sigma = 3\%$. Therefore, we conclude that *i*) the VRE on these sources is measured to be $VRE = 0\%$, with $\sigma \approx 3\%$, and *ii*) the VRE we measured in the Earth albedo ranges between 4 and 10%, with a similar error of $\sigma \approx 3\%$ (Table 3). In the June 17+18 spectrum, we probably have a larger error on the VRE due to the lack of bracketing from Earthshine with Moonlight spectra (Fig. 6 and 7).

These results are in agreement with estimations from models: Des Marais *et al.* (2001) predict a vegetation signature of “2%, maybe larger if a large forested area is in view”. Preliminary estimates gave 5% (Schneider 2000a, Schneider 2000b), while our model described hereafter predicts $\approx 10\%$.

Only green lands observed by different Earth observing satellites show this spectral feature around 700 nm. It is thus legitimate to attribute our global VRE to the terrestrial vegetation. Let us investigate this hypothesis a little further.

Table 4 shows which portion of the Earth is seen from the Moon at the time the observations were done. An important contribution to the Earth albedo comes from clouds. When clouds cover land without vegetation or an ocean area, its albedo adds to the general planet albedo, without suppressing the vegetation contribution, and simply reduces the global VRE. When clouds cover a forest region, the vegetation contribution to the VRE for that Earth region is removed. The global VRE thus obviously depends not only on the global cloud cover, but also on which re-

Table 4. Earth zone viewed from the Moon at the epochs of observations.

Epoch	Viewed zone	Land %	Ocean %
2001/06/17&18	Eur./Asia	50	50
2001/06/24&26	Amer./Pacific	40	60
2001/07/23&24&25	Amer./Pacific	40	60
2001/10/13&14	Eur./Asia	50	50
2001/10/19&21	Amer./Pacific	30	70

gions are covered by clouds during the observation. We have therefore taken cloud cover images available from <http://wwwghcc.msfc.nasa.gov/GOES/globalir.html> for each of our observation dates and have estimated for each run of observation the fraction of land and ocean covered by clouds (Table 5).

One can *a priori* estimate the *VRE* from the albedo and relative surfaces of ocean, land and cloud. Using Eq.6, the *VRE* can be written

$$VRE_{th} = \frac{A_l(r_I - r_R)_l CF_l}{A_o(r_R)_o CF_o + A_l(r_R)_l CF_l + CC(r_R)_c} \quad (8)$$

where $CF_{o,l}$ are the fractional cloud-free areas for ocean and land (from deserts to green forests), respectively, and $A_{o,l}$ are the corresponding areas in view from the Moon. CC is the fractional cloud cover, equal to $CC = (1 - CF_o)A_o + (1 - CF_l)A_l$. Calculating a mean VRE_{th} for Earth with $A_o = 0.7$ and $A_l = 0.3$ (standard Earth ocean/land ratio), $(r_I - r_R)_l = 0.15$ (based on *NDVI* data), $(r_R)_o = 0.05$, $(r_R)_l = 0.25$ and $(r_R)_c = 0.5$ (albedos for ocean, land and cloud), we find, for a mean cloud cover of 50% for all regions, $VRE_{th} \approx 7\%$. We estimate the error on VRE_{th} to be $\pm 3\%$ (cloud cover estimation, seasonal variation of the difference $(r_I - r_R)_l$, vegetation mutual shadow effects). The small contribution of ocean chlorophyll is also neglected here.

The resulting VRE_{th} computed from the Earth phase seen from the Moon (Table 4) and cloud cover (Table 5) are given in Table 6. The denominator of Eq.8 represents an albedo, known to be 0.30 for the Earth (Goode *et al.* 2001), and its value here ranges between 0.30 and 0.34 depending on the date of observation. The VRE_{th} values are in acceptable agreement with preliminary estimates of 5% (Schneider 2000a, Schneider 2000b), but are higher than our observations. This partially comes from terrestrial limb and directionality of the vegetation reflectance effects. The latter are complicated (Privette *et al.* 1995, Hapke *et al.* 1996), due for instance to mutual shadow effects between trees.

5. Conclusion

Although it seems that the Earth’s vegetation signature might be visible as a red edge at 700 nm, it is difficult to

Table 5. Cloud cover estimated for the Earth zone viewed from the Moon at the epochs of observations.

Epoch	Land (1 - CF_l)%	Ocean (1 - CF_o)%
2001/06/17&18 morning	60	50
2001/06/24&26 evening	40	50
2001/07/23&24&25 evening	50	50
2001/10/13&14 morning	50	50
2001/10/19&21 evening	50	50

Table 6. Comparison of the observed and calculated Vegetation Red Edge *VRE*.

Epoch	Viewed zone	VRE_{obs}	VRE_{th}
2001/06/17&18 morning	Eur./Asia	$10 \pm 5\%$	$9 \pm 3\%$
2001/06/24&26 evening	Amer./Pacific	$7 \pm 3\%$	$12 \pm 3\%$
2001/07/23&24&25 evening	Amer./Pacific	$7 \pm 3\%$	$10 \pm 3\%$
2001/10/13&14 morning	Eur./Asia	$4 \pm 3\%$	$12 \pm 3\%$
2001/10/19&21 evening	Amer./Pacific	$7 \pm 3\%$	$7 \pm 3\%$

measure in the Earthshine for two reasons. The first reason is related to its variable amplitude, induced by a variable cloud cover and Earth phase. The second reason is because it is hidden below strong atmospheric bands which need to be removed to access the surface reflectance including the vegetation signature.

For the Earth, our knowledge of different surface reflectivities (deserts, ocean, ice etc) help us to assign the *VRE* of the Earthshine spectrum to terrestrial vegetation. For an exoplanet, a *VRE*-like index might be as difficult to measure as for the Earth due to variable cloud cover of the planet. Even if an extrasolar planet would give a clear *VRE*-like spectral signal, its use as a biosignature would raise some questions because:

1/ For several organisms (such as *Rhodospseudomonas*, Blankenship *et al.* 1995) the “red edge” is not at 700 nm, but at 1100 nm.

2/ Some rocks, like schists, may have a similar spectral feature. For instance, spectra of Mars show a similar spectral feature at 3.5μ , which were erroneously interpreted as vegetation due to their similarity with lichen spectra (Sinton 1957).

We nevertheless believe that, associated with the presence of water (and secondarily oxygen) and correlated with seasonal variations, a vegetation-like spectral feature would provide more insight than simply oxygen on the bio-processes possibly taking place on the planet. But since water, and thus clouds and rain, are essential for the growth of vegetation, extrasolar planets with a very low cloud cover and a corresponding high vegetation index are unlikely, more especially if the planet is seen pole-on, with a bright white polar cover. On the other hand, an extra-

solar planet vegetation surface could be larger than on Earth (like during periods in the paleozoic and mesozoic eras on Earth for example).

One must also note that the measurement of an extrasolar planet VRE will not suffer from the intrinsic difficulty of the same measurement for the Earth through the Earthshine spectrum: The extrasolar planet albedo will simply be given by the ratio of spectra *planet/mother star*. But a model of the exoplanet atmosphere is necessary to be able to remove the absorption bands that may partially hide the vegetation. The detection of a VRE index between 0 and 10% requires a photometric precision better than 3%. Exposure time to achieve this precision with Darwin/TPF on an Earth-like planet at 10 *pc* with a spectral resolution of 25 is of the order of 100 *h* based on recent simulations (Riaud *et al.* 2002).

Finally, the Earth albedo spectral variations study is of interest for global Earth observation. It might provide data on climate change, as broad-band measurements recently showed (Goode *et al.* 2001). We also think that the spectrum of Earthshine might be used for example to monitor the global ozone (with the Chappuis or Huggins bands).

During the submission of the paper, we have been informed of a similar work by Woolf *et al.* (2002).

Acknowledgements. We are grateful to P. François who took test spectra of the Earthshine in 1999. We thank Ph. Gastellu-Etchegorry for providing us Earth observation satellite data. We also thank C. Prigent, L. Beaufort, G. Bellucci, D. Gillet, S. Le Mouélic, A. Labeyrie, J.M. Perrin and referee S. Franck for constructive remarks about this work. F. Valbousquet from *Optique et Vision* lent us some parts for the spectrograph.

References

- Angel J.R.P., Chen A.Y.S., Woolf N.J. 1986, *Nature*, 322, 341
 Angel J.R.P., Woolf N.J. 1997, *ApJ*, 475, 373
 Arcichovsky V.M. 1912, 'Auf der Suche nach Chlorophyll auf den Planeten' in *Annales de l'Institut Polytechnique Don Cesarevitch Alexis a Novotcherkassk*, Vol.1 no 17, 195
 Beichman C., Lindensmith C., Woolf N.J. 1999, *The Terrestrial Planet Finder*, JPL Publication 99-3
 Blankenship R.E., Madigan M.T., Bauer C.E. 1995, *Anoxygenic Photosynthetic Bacteria* (Kluwer Academic Publishing, Dordrecht, The Netherlands)
 BRDF (Special issue on) 2000, *Remote Sensing Reviews*, 18, 83 to 551
 Clark R.N. 1999, *Manual of Remote Sensing* (J. Wiley and Sons, ed. A. Rencz, New-York)
 Coliolo F., Labeyrie A., Schneider J. 2000, in *Proc., Sixth Trieste Conference on Chemical Evolution 'First Steps in the Origin of Life in the Universe'*
 Danjon A. 1928, *Ann. Obs. Strasbourg*, 2, 165
 Des Marais D., Harwit M., Jucks K., Kasting J., Lin D., Lunine J., Schneider J., Seager S., Traub W., Woolf N. 2001, *JPL Publication 01-008*
 Goode P., Qiu J., Yurchyshyn V., Hickey J., Chu M.-C., Kolbe E., Brown C., Koonin S. 2001, *Geophys. Res. Lett.*, 28(9), 1671
 Hapke B., Dimucci D., Nelson R., Smythe W. 1996, *Lunar and Planetary Sci.*, 27, 491
 Kelley K.W. and the Association of Space Explorers 1988, *The Home Planet* (Addison-Wesley Publishing Company, New-York)
 Labeyrie A., Schneider J., Boccaletti A., Riaud P., Moutou C., Abe L., Rabou P. 1999, in *ESA Proc., Darwin and Astronomy*, ESA SP-451, 21
 Labeyrie A. 1999, in *Proc., Planets Outside the Solar System: Theory and Observations*, eds. J.M. Mariotti and D. Alloin, *NATO ASI* 532, 261
 Léger A., Mariotti J.M., Menesson B., Ollivier M., Puget J.L., Rouan D., Schneider J. 1996, *Icarus*, 123, 249
 Liang S., Strahler A.H. 1999, *The Earth Observer*, 11, 27
 Lovelock J. 1975, *Proc. Roy. Soc. London*, B 189, 167
 McLinden C.A., McConnell J.C., Griffioen E., McElroy C.T., Pfister L. 1997 *J. Geophys. Res.*, 102, 18801
 Owen T. 1980, in *Proc., Strategies for the search for life in the universe*, ed. M. Papagiannis, Reidel, 177
 Pieters C.M. 1999, in *Proc., New Views of the Moon II: Understanding the Moon Through the Integration of Diverse Datasets*, September 22-24, 1999, Flagstaff, Arizona, abstract 8025 (<http://www.planetary.brown.edu/pds/AP62231.html>)
 Prigent C., Aires F., Rossow W., Matthews E. 2001, *J. Geophys. Res.*, 106, 20665
 Privette J.L., Myneni R.B., Emery W.J., Pinty B. 1995, *J. Geophys. Res.*, 100, 25497
 Rouse J.W., Haas R.H., Schell J.A., Deering D.W. 1974, *Final Report, Type III, NASA/GSFC, Greenbelt*
 Riaud P., Boccaletti A., Gillet S., Arnold L., Lardiére O., Dejonghe J., Labeyrie A. 2002, *A&A*, to be submitted
 Sagan C., Thompson W.R., Carlson R., Gurnett D., Hord C. 1993, *Nature*, 365, 715
 Schneider J. 2000a, *Exoplanets in A Encyclopaedia of Astronomy and Astrophysics* (Institute of Physics Publishing)
 Schneider J. 2000b, private communication
 Schneider J., Auvergne M., Baglin A., Michel E., Rouan D., Appourchoux T., Barge P., Deleuil M., Vuillemin A., Catala C., Garrido R., Léger A., Weiss W. 1998, in *ASP Conf. Ser. 148*, ed. Charles E. Woodward, J. Michael Shull, and Harley A. Thronson, Jr., 298
 Sinton W. 1957, *ApJ*, 126, 231S
 Tikhoff G.A. 1914, *Mitteilungen der Nikolai-Hauptsternwarte zu Pulkowo*, no 62, Band VI₂, 15
 Tucker C.J. 1979, *Remote Sensing of the Environment*, 8, 127
 Udry S., Mayor M. 2001, in *ESA Proc., First European Workshop on Exo/Astrobiology*, in press
 Woolf N., Smith P., Traub W., Jucks K. 2002, accepted by *ApJ* 20 March 2002
 Wolszczan A., Frail D. 1992, *Nature*, 255, 145